



AFRL-ML-WP-TP-2007-495

**ELECTRIC-FIELD EFFECTS IN ESR SPECTRUM OF
LOW-SPIN CENTER Ni^{3+} IN KTaO_3 CRYSTALS
(PREPRINT)**

D.R. Evans

**Hardened Materials Branch
Survivability and Sensor Materials Division**

APRIL 2007

Approved for public release; distribution unlimited.

See additional restrictions described on inside pages

STINFO COPY

**AIR FORCE RESEARCH LABORATORY
MATERIALS AND MANUFACTURING DIRECTORATE
WRIGHT-PATTERSON AIR FORCE BASE, OH 45433-7750
AIR FORCE MATERIEL COMMAND
UNITED STATES AIR FORCE**

NOTICE AND SIGNATURE PAGE

Using Government drawings, specifications, or other data included in this document for any purpose other than Government procurement does not in any way obligate the U.S. Government. The fact that the Government formulated or supplied the drawings, specifications, or other data does not license the holder or any other person or corporation; or convey any rights or permission to manufacture, use, or sell any patented invention that may relate to them.

This report was cleared for public release by the Air Force Research Laboratory Wright Site (AFRL/WS) Public Affairs Office and is available to the general public, including foreign nationals. Copies may be obtained from the Defense Technical Information Center (DTIC) (<http://www.dtic.mil>).

AFRL-ML-WP-TP-2007-495 HAS BEEN REVIEWED AND IS APPROVED FOR PUBLICATION IN ACCORDANCE WITH ASSIGNED DISTRIBUTION STATEMENT.

*//Signature//

DEAN R. EVANS, Ph.D.
Agile Filters Project
Exploratory Development
Hardened Materials Branch

//Signature//

MARK S. FORTE, Acting Chief
Hardened Materials Branch
Survivability and Sensor Materials Division

//Signature//

TIM J. SCHUMACHER, Chief
Survivability and Sensor Materials Division

This report is published in the interest of scientific and technical information exchange, and its publication does not constitute the Government's approval or disapproval of its ideas or findings.

*Disseminated copies will show “//Signature//” stamped or typed above the signature blocks.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
<p>The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</p>					
1. REPORT DATE (DD-MM-YY) April 2007		2. REPORT TYPE Journal Article Preprint		3. DATES COVERED (From - To)	
4. TITLE AND SUBTITLE ELECTRIC-FIELD EFFECTS IN ESR SPECTRUM OF LOW-SPIN CENTER Ni ³⁺ IN KTaO ₃ CRYSTALS (PREPRINT)				5a. CONTRACT NUMBER In-house	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER 62102F	
6. AUTHOR(S) L.S. Sochava, S.A. Basun, V.E. Bursian, and A.G. Razdobarin (A.F Ioffe Physico-Technical Institute of RAS) D.R. Evans (AFRL/MLPJ)				5d. PROJECT NUMBER 4348	
				5e. TASK NUMBER RG	
				5f. WORK UNIT NUMBER M08R1000	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) A.F Ioffe Physico-Technical Institute of RAS 194021 Saint-Petersburg, Russia				Hardened Materials Branch (AFRL/MLPJ) Survivability and Sensor Materials Division Materials and Manufacturing Directorate Wright-Patterson Air Force Base, OH 45433-7750 Air Force Materiel Command, United States Air Force	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Air Force Research Laboratory Materials and Manufacturing Directorate Wright-Patterson Air Force Base, OH 45433-7750 Air Force Materiel Command United States Air Force				10. SPONSORING/MONITORING AGENCY ACRONYM(S) AFRL/MLPJ	
				11. SPONSORING/MONITORING AGENCY REPORT NUMBER(S) AFRL-ML-WP-TP-2007-495	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited.					
13. SUPPLEMENTARY NOTES Journal article submitted to Physics of the Solid State. The U.S. Government is joint author of this work and has the right to use, modify, reproduce, release, perform, display, or disclose the work. PAO Case Number: AFRL/WS 07-0773, 03 Apr 2007.					
14. ABSTRACT Electric-field effects in ESR spectrum of low-spin (S=1/2) tetragonal center Ni ³⁺ was studied in KTaO ₃ single crystals. Orientation of the centers as well as splitting of the resonance lines was found resulting from the external E-field interaction with the electric dipole moment of the center. The value of the dipole moment was determined to be p=103D=21.4 eÅ. An analysis of the experimental results allowed us to make a reasonable choice of microscopic models for two nickel centers in KTaO ₃ .					
15. SUBJECT TERMS KTaO ₃ crystals, ESR Spectrum, Ni ³⁺					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT: SAR	18. NUMBER OF PAGES 10	19a. NAME OF RESPONSIBLE PERSON (Monitor) Dean R. Evans
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified			19b. TELEPHONE NUMBER (Include Area Code) N/A

Electric-field effects in ESR spectrum of low-spin center Ni^{3+} in KTaO_3 crystals.

L.S. Sochava^a, S.A. Basun^a, V.E. Bursian^a, A.G. Razdobarin^a, D.R. Evans^b

^a *A.F. Ioffe Physico-Technical Institute of RAS, 194021 Saint-Petersburg, Russia*

^b *Air Force Research Laboratory, Materials and Manufacturing Directorate, Wright-Patterson Air Force Base, Ohio, 45433, USA*

Electric-field effects in ESR spectrum of low-spin ($S=1/2$) tetragonal center Ni^{3+} was studied in KTaO_3 single crystals. Orientation of the centers as well as splitting of the resonance lines was found resulting from the external E-field interaction with the electric dipole moment of the center. The value of the dipole moment was determined to be $p=103D=21.4 \text{ eÅ}$. An analysis of the experimental results allowed us to make a reasonable choice of microscopic models for two nickel centers in KTaO_3 .

This study was supported by the Russian Foundation for Basic Research (project no. 06-02-17274) and the European Office of Aerospace Research and Development (CRDF grant no. RPO-1385-ST-03).

1. Introduction

Iron-group impurities in KTaO_3 are known to enter substitutionally on either tantalum or potassium site. Besides, these 3-d ions can have different defects in a close proximity, thus forming a wide variety of defect centers in potassium tantalate.

In late sixties, two tetragonal Ni^{3+} centers were found in KTaO_3 [1]; recently we have observed a nickel center of orthorhombic symmetry [10]. Unlike two Fe^{3+} centers found in KTaO_3 [14], tetragonal Ni^{3+} centers can have two different spin values – low-spin ($S=1/2$) and high-spin ($S=3/2$) depending on the type of cation nickel substitutes for (Ta site has an octahedral surrounding whereas K site has an icosahedral surrounding).

However, the exact positions of each of the two centers remain unclear. Most likely, this problem can not be solved with the use of only Hund's rule, without knowledge of three characteristic energies: crystal field splitting of the levels $10Dq$, spin-orbit coupling, and exchange interaction.

The origin of tetragonal distortion of the crystal field is another open question. The authors of [1] suggested a Jahn-Teller nature of tetragonal distortion, others believe it to be rather a

neighboring oxygen vacancy [3], potassium vacancy [2], interstitial oxygen, or a noncentral position of nickel ion [2].

This paper presents the study of the electric-field effects in the ESR spectrum of the low-spin ($S=1/2$) center Ni^{3+} in KTaO_3 . An analysis of the experimental results allowed us to choose a microscopic model for the center.

2. Experiment technique

The experiments were performed on the samples of potassium tantalate grown in the Laboratory of Crystal Growth of Osnabrück University, Germany. Rectangular samples with the edges aligned with $\langle 100 \rangle$ and measuring typically $0.9 \times 4 \times 6 \text{ mm}$ were cut from a single-crystal boule. Aquadag-paste electrodes were deposited on $4 \times 6\text{-mm}$ faces. The sample with electrodes was immersed in an insulating paste. This made it possible to apply a high voltage (up to 10 kV) across the sample, not only in liquid but also in gaseous nitrogen. The ESR spectra were measured on a modified 3-cm Radiopan SE/X2544 spectrometer with 100-kHz magnetic field modulation.

ESR signals from both low-spin ($S=1/2$) and high-spin ($S=3/2$) centers are observed in as-grown

KTaO₃ samples exposed to UV light at 78 K. Subsequent irradiation of the sample with red light increases concentration of high-spin centers and at the same time decreases that of low-spin centers.




Fig. 1. Direction of the electric and magnetic fields

3. Linear dependence of line splitting on electric field

An external electric field up to 105 kV/cm was applied at liquid nitrogen temperature along one of the cubic axes of the KTaO₃ crystal (Fig.1). In this layout, for the centers with the axis parallel to x, an electric-field effect manifests itself in a partially resolved line splitting (Fig. 2).




Fig. 2. Splitting in the external electric field of the ESR line of the low-spin-Ni³⁺ center oriented along x, E || y, E = 105 kV/cm, T = 78 K, ν = 9.4 GHz. Circles — the values extracted from the experiment as explained in the text, line — linear fit.

The splitting appears as soon as the field is applied, and there are no further changes in the splitting value or in the intensities of the components. This means that there are two states of the center, which are unresolved in the absence of the electric field.

At lower values of the applied field the effect is observed as a broadening of the line that initially has a width of 6.5 Oe. In order to extract the values

of line splitting, we made a fit of the experimental line shape with the zero-field line split in two halves by a variable parameter. Fig. 2 shows the splitting parameter that gives the best match, which yields a linear dependence.

4. Alignment of Ni Centers in the Electric Field

Linearity of the electric-field effect in the ESR spectrum of the discussed center is evidence for a non-inversion symmetry in the site with substituent nickel. In other words, the discussed center must have an electric dipole moment.

To be oriented by an external electric field, the dipole center must have a possibility to undergo spontaneous (thermal) reorientations at the temperature of the experiment. These reorientations are not observed at T=78 K, but they are detectable at higher temperatures. A distinct alignment of nickel centers was observed between 120-130 K.

By varying a temperature within this range, we have determined a value of the electric dipole moment. The ratio of intensities of the lines related to differently oriented centers was measured in a two-step procedure:

- an E-field was applied to the sample along one of the cubic axes at 120-130 K. In this temperature interval, the equilibrium between the centers aligned along and perpendicular to the direction of E-field is established rather quickly (2-3 minutes at 125 K).
- Subsequently, the temperature was rapidly (in a 5-10 seconds) decreased to 100 K (the characteristic reorientation time at 100 K is about few days), the E-field was switched off, and ESR spectrum was recorded. Recording the spectra in zero E-field allows for the avoidance of a problem with line shape distortion by external field.

The ratio n_y/n_x , where n_y and n_x are intensities of the resonance lines of the Y and X centers (E-field was parallel to Y axis) was plotted in coordinates $\text{arch}(n_y/n_x)$ vs. E/kT (Fig.3). The dependence appears linear with a good accuracy. The data on Fig. 3 were obtained by varying the external E-field from 0 to 74 kV/cm and the temperature from 122 to 130 K. The slope ratio of the line is equal to the electric dipole moment of the center p :

$$p = (21.4 \pm 2.0) \text{ eÅ} = (103 \pm 10) \text{ D}$$




Fig. 3. $\text{arch}(n_y/n_x)$ as a function of E/kT . The values were obtained from the measured ratios of the concentrations of the centers oriented along y and x axes at different temperatures and external fields.

5. Line splitting dependence on the orientation of electric and magnetic fields

Fig. 4 shows the angle dependence of resonance line splitting for the centers oriented along x axis, when electric field is *perpendicular* to the axis of the center ($E \parallel y$) and the magnetic field rotates in the (xz) plane (Fig. 1). The splitting equals zero for the directions $H \parallel x$, $H \parallel y$, and $H \parallel z$; it reaches a maximum when H lies in (xz) plane at $\sim 45^\circ$ to the x-axis.




Fig. 4. Correlation between the angular dependence of the splitting in the electric field and that of the first derivative of the resonant magnetic field. Circles – splittings in the electric field (left axis), solid curve – first derivative of resonant magnetic field (right axis), the centers are oriented along x-axis, $E = 105$ kV/cm.

To the best of our knowledge, a similar manifestation of electric-field effect in ESR was observed only once [9] on the $\text{Fe}^{3+}_{\text{K-O}}$ center (in potassium tantalite also).

It is explained as follows: in the electric field perpendicular to the axis of the center, the dipole center slightly rotates by a small angle. Consequently, the angle between the axis of the center and the direction of the magnetic field changes. In the layout shown in Fig. 1 for the center with dipole moment along +x, this angle slightly decreases, whereas for the center with dipole moment along -x, it slightly increases; therefore, the shifts are opposite and the line splits. The value of the splitting must be directly correlated with the first derivative of the resonant magnetic field as a function of angle in this plane, which, indeed, takes place as one can see on Fig. 4.

In addition to the effect described in [9], we have observed also a splitting of the resonance lines (smaller in magnitude than one described above) for the centers *parallel* to the external electric field. Although we failed to obtain a presentable angle dependence $\delta H(H)$ in this layout, the presence of the effect is beyond any doubts.

6. Discussion on the microscopic structure of the center

6.1 As noted in the introduction, two tetragonal ESR spectra are observed in $\text{KTaO}_3:\text{Ni}$ and attributed to Ni^{3+} ions with different values of effective spin. To clarify the position of each of the two Ni^{3+} ions in the lattice of potassium tantalate we compared electric-field effects observed on Ni^{3+} ions with those found for Fe^{3+} in KTaO_3 . As mentioned above, the main features of electric-field effect for Ni^{3+} centers perpendicular to E-field are similar to the results reported for Fe^{3+} ions in potassium site in KTaO_3 ($\text{Fe}^{3+}_{\text{K-O}}$) [9]. Authors of [9] suggested the mechanism for the effect to be a rotation of the dipole center, by a small angle, in the external electric field. It is important that the ionic radii of Ni^{3+} and Fe^{3+} ions are both much smaller than that of K^+ ($\sim 1.4 \text{ \AA}$), which makes it easier for the centers to deviate in an external field. We believe this is a strong argument for the potassium site to be occupied by the low-spin nickel ion. On the other hand, the electric field does not split resonance lines in the ESR spectrum of the high-spin ($S=3/2$) Ni^{3+} center. Absence of the electric field effect was also reported for the $\text{Fe}^{3+}_{\text{Ta}}$ center, iron in the tantalum site, [13] and this justifies our assignment.

6.2 Another issue regarding the microscopic structure of nickel centers that remained unclear up to date is the origin of tetragonal symmetry of the crystal field. The linearity of the electric-field effect reported in this paper allows us to disregard the idea of Jahn-Teller nature of tetragonal distortion of the low-spin nickel center in potassium tantalite as given earlier [1].

Furthermore, the dipole moment of the center is very unlikely to arise from a noncentral position of the nickel ion because of a relatively large value of the dipole moment ($p=21.4$ eÅ). On the other hand, the value found for p differs by only 30% from that of the $Fe^{3+}_{K-O_i}$ center in $KTaO_3$ [13]; the structure of the latter was reliably established in [14] by means of ENDOR. This allows us to suggest that the low-symmetry center of nickel has a similar structure, *i.e.* Ni^{3+} in potassium site next to an interstitial oxygen ($Ni^{3+}_{K-O_i}$).

Acknowledgments

The authors are grateful to Dr. H. Hesse and Prof. S. E. Kapphan for providing high-quality $KTaO_3:Ni$ crystals, to Prof. V. S. Vikhnin for fruitful discussions, and to Prof. A. A. Kaplyanskii for interest in the work.

References

1. D.M. Hannon. *Phys. Rev.* **164**, 366 (1967).
2. Y. Akishige and K. Ohi. *Jap. J. Appl. Phys.* **19**, 1633 (1980).
3. M.M. Abraham, L.A. Boatner, D.N. Olson, U.T. Höchli. *J. Chem. Phys.* **61**, 2528 (1984).
4. D.M. Hannon. *phys. status solidi B* **43**, K21 (1971).
5. A.G. Badalyan, P.G. Baranov, D.V. Azamat, V.A. Trepakov, C.B. Azzoni, P. Galinetto, M.C. Mozzati, L. Jastrabik, J. Rosa. *J. Phys.: Condens. Matter* **14**, 6855 (2002).
6. S. A. Basun, L. S. Sochava, V. E. Bursian, H. Hesse, S. Kapphan, V. S. Vikhnin. *Proc. of the XIII Internat. Conf. on Defects in Insulating Materials*, Wake Forest University, USA, July 15-19, 1996, in *Materials Science Forum*, Trans Tech Publications, Switzerland **239-241**, 345 (1997).
7. S.A. Basun, V.E. Bursian, A.G. Razdobarin, L.S. Sochava. *Physics of the Solid State* **43**, 1059 (2001).
8. S.A. Basun, A.G. Razdobarin, L.S. Sochava, D.R. Evans. *Physics of the Solid State* **46**, 258 (2004).
9. I.N. Geifman, M.D. Glinchuk, B.K. Krulikovskii. *Zh. Exp. Teor. Fiz.* **75**, 1468 (1978).
10. L.S. Sochava, V.E. Bursian, A.G. Razdobarin, S.A. Basun, D.R. Evans, S.E. Kapphan. *Physics of the Solid State* **49**, 443 (2007).
11. Y. Akishige, Y. Yamamoto, K. Ohi. *Jap. J. Appl. Phys.* **21**, 1037 (1982).
12. S.Y. Wu, H.N. Dong, W.H. Wei. *Z.Naturforsch.* **59a**, 203 (2004).
13. V.V.Laguta, M.D.Glinchuk, A.A.Karmazin, I.P.Bykov. *Physics of the Solid State* **27**, 162 (1985).
14. I.P.Bykov, M.D.Glinchuk, A.A.Karmazin, V.V.Laguta. *Physics of the Solid State* **25**, 3586 (1983).